The NUTEV DIS $sin^2\theta_W$ Anomaly or Beware of Mass Singularities (ln(M/m_f) terms)

William J. Marciano
December 11, 2020
Lunch Discussion



Based on: Oldies but Goodies

WJM & A. Sirlin (1973) *electron-muon universality*

WJM (1975) Dim. Reg. of mass singularities

D. Albert, WJM, Z. Parsa, D. Wyler (1980) W & Z Decays

*A. Sirlin and WJM (1981) Neutrino DIS & $\sin^2 \theta_W$

More Recent

A. Czarnecki, WJM, G.P. Lepage (2000) *Muonium Decay*


2020 Nobel Prize Connection

Black Holes and BNL

Precision EW Parameters (status):

Quantity	<u>2008 Value</u>		<u>2020 Value</u>	<u>Commen</u> t	
$\alpha^{\text{-1}}$	137.035999084	(51)	137.035999046(27)	$\alpha^{-1}(Cs)$	
			137.035999150(34)	$lpha^{\text{-1}}(a_e)$	
			137.035999206(11)	$lpha^{\text{1}}$ (Rb) 5.5 σ tension	New!
G_{μ}	1.16637(1)x10 ⁻⁵ G	eV ⁻²	1.1663787(6)x10 ⁻⁵ GeV ⁻²	τ_{μ^+} PSI MuLAN 2010	
m_{z}	91 .1875(21)GeV	,	91.1876(21)GeV	-	
*m _t	171.4(2.1)GeV	\rightarrow	172.9(0.4)GeV	FNAL/LHC	
$*m_H$	>114GeV	\rightarrow	125.10(0.14)GeV	LHC	
m_{W}	80.410(32)GeV	\rightarrow	80.379(12)GeV	LEP2/FNAL/LHC	
			80.359(3)GeV	Global Fit Value	
$sin^2\theta_W(m_Z)$	0.23070(26)		0.23070(26)	SLAC A _{LR}	
$sin^2\theta_w(m_z)$	0.23193(29)		0.23193(29)	CERN A _{FB} (bb)	
(3 sign	na difference?)				
$\sin^2\theta_W (m_Z)_{ave} 0.23125(16)$		0.23119(14)	Z Pole Ave.		
			<u>0.23120(6)</u>	Global EW Fit	

Comparison of recent α determinations from Nature (2020) 5.5 sigma difference between α (Rb) and α (Cs) $\Delta a_e = a_e^{\text{exp}} - a_e^{\text{SM}} \text{ changes sign!}$



Comment on New $\alpha(Rb)$ value

$$\Delta a_e = a_e^{exp} - a_e^{SM} = -87(28)_{exp}(23)_{\alpha}(2)_{th} \times 10^{-14}$$
 for Cs $\Delta a_e = a_e^{exp} - a_e^{SM} = +47(28)_{exp}(9)_{\alpha}(2)_{th} \times 10^{-14}$ for Rb sign change $\Delta a_{\mu} = a_{\mu}^{exp} - a_{\mu}^{SM} = +261(40)(26)\times 10^{-11}$ $\frac{\Delta a_{\mu}}{\Delta a_e} \sim 5553$ if m_{μ}^2/m_e^2 scaling holds, e/μ universality, $\Delta a_e = a_e^{exp} - a_e^{SM} \sim 6\times 10^{-14}$ expected

Accommodate with a light Scalar But first overcome Rb/Cs Difference


Best Off Z Resonance Measurements of $\sin^2\theta_W(m_Z)_{MS}$ (Not Yet Competitive with Z Pole)

Reaction	$\sin^2\theta_W(m_Z)_{MS}$	<q></q>
Cs APV	0.2283(20)	2.5MeV
E158 ee	0.2329(13)	160MeV
Q _{weak} ep	0.2310(11)	160MeV
Average	0.23127(77)	Good Agreement
Z Pole	0.23119(14)	
*NuTeV ν _μ N	0.23560(160)	10 GeV


NuTeV high by about 2-3 sigma using R_{ν}^{exp} = σ_{NC}/σ_{CC} =0.3916(7)

BSM Unlikely? R_{ν} incomplete Radiative Corrections? Likely!

Running of the Weak Mixing Angle



Running of $sin^2\vartheta_W(Q)$ & Future Measurements



Radiative corrections to $R_{\overline{\nu}}$ (A. Sirlin &WJM 1981)

$$R_{\nu}^{exp} \equiv \sigma(\nu_{\mu}N \rightarrow \nu_{\mu}X)/\sigma(\nu_{\mu}N \rightarrow \mu X') = 0.3916(7) \text{ NuTeV}.$$
 (2002)

Radiative Corrections 1 loop + Bremsstrahlung (Monte Carlo)

Experiment:
$$\sin^2 \theta_W \equiv 1 - \frac{m_W^2}{m_Z^2} = 0.2277(16) \ vs \ \text{SM global fit} \rightarrow \underline{\textbf{0.2234(1)}}$$

Main RC from W γ Box diagram increases CC cross-section by $\frac{\alpha}{\pi} (ln \frac{m_Z^2}{2Q^2} + 2) \sim 1.8\%$

Suggests R_{ν}^{SM} needs additional correction by $\sim 0.9\%$ to agree with Global average

<u>Mass Singularities theorem (1959)</u> No large $O(\frac{\alpha}{2\pi} ln(\frac{2mNE}{m_{\mu}^2})) \sim 1\%$


corrections for $\nu_{\mu}N \rightarrow \mu X'$ fully inclusive cross-section over all E_{μ} .

However, <u>cuts on Bremsstrahlung induce mass singularities</u>. Increase CC cross-section by about +0.8-0.9%. Just what is needed for agreement with SM. <u>Already included??</u>

Predicted by A. Sirlin &WJM in 1981! Generic +0.7%

Recently gifted to NuTeV Collaboration for consideration. The Jury is still out.

Box Diagram Loop Induced RC



Example of cuts: e/μ Universality Breakdown in $W \rightarrow l\nu(\gamma)$ decays ($l=e,\mu$)

Dimensional Regularization of both infrared div. & lepton mass sing. $m_{\gamma}=0$ $m_{l}=0$ Compare with $m_{l}\neq 0$ calculation, neglecting $O(\frac{m_{l}^{2}}{m_{W}^{2}})$ effects taking $n\rightarrow 4$

Inclusive Decay Rate=
$$g^2 \frac{m_W}{48\pi} (1 - \frac{\alpha}{2\pi} (\frac{2\pi}{3} - \frac{77}{12})) \rightarrow \frac{\Gamma(W \rightarrow e\nu(\gamma))}{\Gamma(W \rightarrow \mu\nu(\gamma))} = 1 + O(\frac{m_l^2}{m_W^2})$$

Exclusive
$$E_{\text{max}} - \Delta E \le E \le E_{\text{max}} \frac{\Gamma(W \to e \nu(\gamma), \Delta E)}{\Gamma(W \to \mu \nu(\gamma), \Delta E)} = 1 - \frac{\alpha}{2\pi} [(4 \ln \frac{m_W}{2\Delta E} - 3) \ln \frac{m_\mu}{me}]$$

Exponentiate
$$=\left(\frac{2\Delta E}{mW}\right)^{0.0248} (1.018)$$

= 0.93 for $\Delta E = 1$ GeV

Cuts can affect Universality tests

eg. B \rightarrow K $l^+ l^-$ New Physics? EXP?

 e/μ detection differences

Precision Muon (μ^+) Lifetime $\rightarrow GF$

MuLAN experiment at PSI:

 τ_{u+} =2.1969803(22)x10⁻⁶sec MuLAN 2010

(Most precise lifetime measurement ever!)

Lowered Previous World Average error by 1/20!

$$\tau_{\mu}^{-1} = \Gamma(\mu^+ \rightarrow e^+ \nu_e \nu_{\mu}(\gamma)) = G_{\mu}^2 m_{\mu}^5 f(m_e^2/m_{\mu}^2) [1 + RC]/192 \pi^3$$

RC = $\alpha/2\pi(25/4-\pi^2)(1+\alpha/\pi[2/3ln(m_{\mu}/m_e)-3.7)...]$ Fermi Th. Other SM and "New Physics" radiative corrections absorbed into G_{μ} . Eg. 4th generation, Technicolor, W*...

 $G_F \equiv G_{\mu} = 1.1663787(6) \times 10^{-5} GeV^{-2}$ precise & important normalization

Loop and Tree Level Corrections to Muon Decay

+ . .

Lessons Learned From Muonium

in Vernon Hughes Memorial Symposium 2003 based on Czarnecki, Lepage & Marciano PRD61 (2000)

MuLan μ^+ Lifetime 1ppm measurement of $\tau_{\mu} \to G_{\mu} \to G_{Fermi}$ Stopped μ^+ in matter forms muonium $M = \mu^+ e^-$ state

Phase space reduction cancels Final State Interaction!

Spectrum Changes $O(\alpha^2 m_e/m_\mu)$ from each

Cancel due to Lorentz & Gauge Invariance

Bound State Ward Identity

 $\tau_{\text{M}} = \tau_{\mu} (1 + \frac{1}{2}\alpha^2 m_e^2 / m_{\mu}^2 + \text{higher order})$

leading correction time dilation≈ O(10-9)

Muon lifetime G_{μ} should not be source of V_{ud} unitarity Problem

 $V_{ud} = 0.97370(14)$: $|V_{ud}|^2 + |V_{us}|^2 + |V_{ub}|^2 = 0.9985(5)$ 3 sigma tension

Phase Space Suppression cancels FSI e⁺e⁻ (many examples)


Similar to Cancellation of Mass Singularities: Kinoshita & Sirlin (1959)

- Muonium $\pm 5\alpha^2 m_e/m_\mu$ linear contributions cancel
- General bound state theorem based on gauge invariance (QED and QCD)
- 1952 The Effect of Atomic Binding on Nuclear Reaction Energies Nuclear Beta Decay <u>Serber & Snyder</u> (first use of theorem)
- 1960 Muon capture Uberall (important potentially large effect)
- Bottom Decays (lifetimes) OPE Proof no 1/m_b effects
- Superallowed Nuclear Beta Decays Complicated at 0.1%

Possible $O(Z^{4/3}\alpha^2m_e/M)$ source of discrepancy?

Future Work

Inventors of Strong Focusing at BNL



Courant Livingston Snyder

The Legacy Of Hartland Sweet Snyder (A Human Interest Story)

"The Effect of Atomic Binding on Nuclear Reaction Energies" by R. Serber and H.S. Snyder PR87(1952). 10 citations (I was last in 2004))


"The strong-focusing synchrotron: A new high-energy accelerator" by Courant, Livingston & Snyder PR88(1952). *135 Citations*

"Quantized space-time" H.S. Snyder PR71(1947)

1751 citations (100 in 2019!)

"On continued gravitation contraction" PR56(1939)

by J. Oppenheimer and H. Snyder (Classical Black Holes Exist!) 876 citations



Snyder and Oppenheimer (1939) (Ahead of its Time)

Roger Penrose Hartland "sweet" Snyder Robert Oppenheimer

Maurice Goldhaber and Ernest Courant BNL Centenarians

